

Wellness Kiosk
Reference

PREMIUM POSTER
11. 5" X 19 "

Main message
displayed here

ENTIRE KIOSK
HEIGHT
5' 5"

FRONT MAGNET
11.25" X 32 "

Front magnet will
contain ISI

PI BOOKLET (US-103748)
[4.25" x 11" folded]
[8.5" x 11" open.]

Note: Please note the
dimensions are in variable
brackets, as PPS is still
determining the exact size
while they lay out the PI and
have the vendor review the
dummy sample.

CVS Brochure (US-102594)

SIDE MAGNET
11.25" X 32 "

Side magnets are
identical

Basket of Ribbons
Image is FPO and does
not reflect what
the actual basket will
look like.

RIBBONS
NOTE: Ribbons
will be white
and be housed
in a basket.



Example of ribbons

Visual does not
reflect the color, as
the ribbons will be
white for Lung Cancer
Awareness Month.

Understanding lung cancer

For adults with EGFR+ non-small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic)



EGFR+=epidermal growth factor receptor mutation-positive.

IMPORTANT PRODUCT INFORMATION

TAGRISSO is a prescription oral treatment used as a first treatment for metastatic non-small cell lung cancer (NSCLC) with certain abnormal EGFR gene(s). TAGRISSO may cause serious side effects, including severe lung problems, heart, eye and skin problems, inflammation of the blood vessels in your skin, and blood and bone marrow problems. Some may lead to death. Tell your healthcare provider right away if you experience trouble breathing, cough, fever, pounding or racing heart, swelling of your ankles and feet, feeling faint, eye problems, skin reactions, including severe blistering or peeling; blood vessel problems, including purple spots or redness of your skin or large hives that do not go away within 24 hours and look bruised, or symptoms of blood and bone marrow problems, including new fever or fever that does not go away ($\geq 100.4^{\circ}\text{F}$), easy bruising or bleeding that will not stop, unusually pale skin, infection, tiredness, or weakness.

Please see full Important Safety Information below and accompanying complete Prescribing Information, including Patient Information for TAGRISSO.



Learn why screening is important



Learn more about lung cancer

These QR codes direct you to third-party websites over which the retailer has no control.

TAGRISSO *I can*

NOVEMBER IS
LUNG CANCER
AWARENESS MONTH



TAGRISSO is a registered trademark of the AstraZeneca group of companies.
©2025 AstraZeneca. All rights reserved. US-102593 8/25

Sponsored by AstraZeneca.

IMPORTANT SAFETY INFORMATION

TAGRISSO may cause serious side effects, including:

- **Lung problems.** TAGRISSO may cause severe lung problems that may lead to death. Symptoms may be similar to those symptoms from lung cancer. Tell your healthcare provider right away if you have any new or worsening lung symptoms, including trouble breathing, shortness of breath, cough, or fever
- **Heart problems, including heart failure.** TAGRISSO may cause heart problems that may lead to death. Your healthcare provider should check your heart function before you start taking TAGRISSO and during treatment as needed. Tell your healthcare provider right away if you have any of the following signs and symptoms of a heart problem: feeling like your heart is pounding or racing, shortness of breath, swelling of your ankles and feet, dizziness, feeling lightheaded, or feeling faint
- **Eye problems.** TAGRISSO may cause eye problems. Tell your healthcare provider right away if you have symptoms of eye problems which may include watery eyes, sensitivity to light, eye pain, eye redness, or vision changes. Your healthcare provider may send you to see an eye specialist (ophthalmologist) if you get eye problems with TAGRISSO
- **Skin problems.** TAGRISSO may cause skin problems. Tell your healthcare provider right away if you develop skin reactions that look like rings (target lesions), severe blistering or peeling of the skin
- **Inflammation of the blood vessels in your skin.** TAGRISSO may cause blood vessel problems in your skin. Tell your healthcare provider right away if you develop purple spots or redness of the skin that does not fade in color when pressed (non-blanching) on your lower arms, lower legs, or buttocks or large hives on the main part of your body (trunk) that do not go away within 24 hours and look bruised
- **Blood and bone marrow problems.** TAGRISSO may cause a condition where your bone marrow cannot make enough new blood cells (aplastic anemia), and which may lead to death. Your healthcare provider will monitor your blood cell counts before you start and during treatment with TAGRISSO. **Tell your healthcare provider right away** if you develop any signs and symptoms of blood and bone marrow problems, including a new fever or fever that does not go away (≥100.4 °F), easy bruising or bleeding that will not stop, unusually pale skin, infection, tiredness, or weakness

Before taking TAGRISSO, tell your healthcare provider about all of your medical conditions, including if you:

- have lung or breathing problems other than your lung cancer
- have heart problems, including a condition called long QTc syndrome
- have problems with your electrolytes, such as sodium, potassium, calcium or magnesium
- have a history of eye problems
- are pregnant or plan to become pregnant. TAGRISSO can harm your unborn baby. Tell your healthcare provider right away if you become pregnant during treatment with TAGRISSO or think you may be pregnant
 - **Females** who are able to become pregnant should have a pregnancy test before starting treatment with TAGRISSO. You should use effective birth control (contraception) during treatment with TAGRISSO and for 6 weeks after the last dose of TAGRISSO
 - **Males** who have female partners that are able to become pregnant should use effective birth control during treatment with TAGRISSO and for 4 months after the last dose of TAGRISSO
- females who are breastfeeding or plan to breastfeed. It is not known if TAGRISSO passes into your breast milk. Do not breastfeed during treatment with TAGRISSO and for 2 weeks after your last dose of TAGRISSO. Talk to your healthcare provider about the best way to feed your baby during this time

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, or herbal supplements. Especially tell your healthcare provider if you take a heart or blood pressure medicine

The most common side effects of TAGRISSO when given alone include:

- low white blood cell counts
- low platelet counts
- low red blood cell counts (anemia)
- diarrhea
- rash
- muscle, bone, or joint pain
- changes in your nails, including: redness, tenderness, pain, inflammation, brittleness, separation from the nailbed, and shedding of nail
- dry skin
- mouth sores
- tiredness

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of TAGRISSO. For more information, ask your healthcare provider or pharmacist.

What is TAGRISSO?

TAGRISSO is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC) that has certain abnormal epidermal growth factor receptor (EGFR) gene(s):

- as your first treatment when your lung cancer has spread to other parts of the body (metastatic)

Your healthcare provider will perform a test to make sure that TAGRISSO is right for you.

It is not known if TAGRISSO is safe and effective in children.

Please see the accompanying complete Prescribing Information, including Patient Information for TAGRISSO.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.





Take the first step

Take a brochure
to learn more



TAGRISSO[®]
osimertinib